

Developmental Pre-Feeding Checklist: A Global Approach

Month Supine with the head slightly elevated; or prone; or reclining it an angle of less than 45 degrees; or sidelying is fed only liquids from the bottle or breast Cakes 2 to 6 ounces of liquid ber feeding, at 6 or more eedings per day	Present- Spontaneous		Not Present
	.ES	Inc.	
Supine with the head slightly elevated; or prone; or reclining t an angle of less than 45 degrees; or sidelying s fed only liquids from the pottle or breast Cakes 2 to 6 ounces of liquid per feeding, at 6 or more eedings per day	.ES	Inc.	
s fed only liquids from the oottle or breast Takes 2 to 6 ounces of liquid per feeding, at 6 or more eedings per day	, ES) ¹	
Cakes 2 to 6 ounces of liquid per feeding, at 6 or more seedings per day	\cap'^*		
ceangs per any	2R		
Jses a suckling or sucking pattern with the bottle or breast. Loses ome liquid during sucking.	5		
wallows thin liquids with a uckle-swallow pattern longue nay protrude slightly through he lips with an extension- etraction movement			
equences two or more sucks			
arely drools because of minimal aliva production			
3 Months			<u> </u>
s fed in a supported semi-sitting position, reclining at an angle of 45 to 90 degrees			
wuunhe ie in waa	vallows thin liquids with a ackle-swallow pattern. Tongue ay protrude slightly through e lips with an extension- traction movement equences two or more sucks on the breast or bottle before ausing to breathe or swallow arely drools because of minimal liva production Months fed in a supported semi-sitting position, reclining at an angle	vallows thin liquids with a ackle-swallow pattern. longue ay protrude slightly through the lips with an extension- traction movement equences two or more sucks on the breast or bottle before ausing to breathe or swallow arely drools because of minimal liva production Months fed in a supported semi-sitting position, reclining at an angle	vallows thin liquids with a ackle-swallow pattern. Tongue ay protrude slightly through e lips with an extension- traction movement equences two or more sucks on the breast or bottle before ausing to breathe or swallow arely drools because of minimal liva production Months fed in a supported semi-sitting position, reclining at an angle