

References

- American Speech-Language-Hearing Association (ASHA). (2004). *Preferred practice patterns for the profession of speech-language pathology*. Retrieved May 3, 2010, from www.asha.org/docs/pdf/PP2004-00191.pdf
- American Speech-Language-Hearing Association (ASHA). (2005). *Roles and responsibilities of speech-language pathologists in diagnosis, assessment, and treatment of individuals with cognitive-communication disorders* [Position Statement]. Retrieved May 3, 2010, from www.asha.org/docs/pdf/PS2005-00110.pdf
- Cicerone, K., Dahlberg, C., Kalmar, K., Langenbahn, D., Malec, J., Bergquist, T., . . . Morse, P.A. (2000). Evidence-based cognitive rehabilitation: Recommendations for clinical practice. *Archives of Physical Medicine & Rehabilitation*, *81*(12), 1596-1615.
- Kennedy, M.R., & Coelho, C. (2005). Self-regulation after traumatic brain injury: A framework for intervention of memory and problem solving. *Seminars in Speech and Language*, *26*, 242-255.
- Shum, D., Gill, H., Banks, M., Maujean, A., Griffin, J., & Ward, H. (2009). Planning ability following moderate to severe traumatic brain injury: Performance on a 4-disk version of the tower of London. *Brain Impairment*, *10*(3), 320-324.