

REFERENCES

- Hargrove, P., Anderson, A., & Jones, J. (2009). A critical review of interventions targeting prosody. *International Journal of Speech-Language Pathology, 11*(4), 298–304.
- Odell, K.H., & Shriberg, L.D. (2001). Prosody-voice characteristics of children and adults with apraxia of speech. *Clinical Linguistics and Phonetics, 15*(4), 275–307.
- Paul, R., Bianchi, N., Augustyn, A., Klin, A., & Volkmar, F. (2008). Production of syllable stress in speakers with Autism Spectrum Disorders. *Research in Autism Spectrum Disorders, 2*(1), 110-124.
- Peppé, S. (2011) Assessment of prosodic ability in atypical populations, with special reference to high-functioning autism. In V. Stojanovik & J. Setter (Eds.), *Speech prosody in atypical populations: Assessment and remediation*. Guildford, U.K.: J & R Press Ltd.
- Ramig, L.O., Countryman, S., Thompson, L.L., & Horii, Y. (1995). Comparison of two forms of intensive speech treatment for Parkinson disease. *Journal of Speech and Hearing Research, 38*, 1232–1251.
- Shriberg, L.D., Paul, R., Black, L.M., & van Santen, J.P. (2011). The hypothesis of apraxia of speech in children with autism spectrum disorder. *Journal of Autism and Developmental Disorders, 41*(4), 405-426. doi: 10.1007/s10803-010-1117-5
- Sparks, R.W., & Holland, A.L. (1976). Method: Melodic intonation therapy for aphasia. *Journal of Speech and Hearing Disorders, 41*, 287–297.
- Strand, E.A., Stoeckel, R., & Baas, B. (2006). Treatment of severe childhood apraxia of speech: A treatment efficacy study. *Journal of Medical Speech-Language Pathology, 14*(4), 297–307.
- Wells, B., Peppé, S., & Goulandris, N. (2004). Intonation development from five to thirteen. *Journal of Child Language, 31*(4), 749–778.